

Uploading 09769450.str

L1 STRUCTURE UPLOADED

=> s sss sam l1

SAMPLE SEARCH INITIATED 14:42:19 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 14361 TO ITERATE

7.0% PROCESSED 1000 ITERATIONS 50 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 280051 TO 294389
PROJECTED ANSWERS: 73816 TO 81282

L2 50 SEA SSS SAM L1

=>

Uploading 09769450b.str

L3 STRUCTURE UPLOADED

=> s sss sam l3

SAMPLE SEARCH INITIATED 14:47:47 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 592 TO ITERATE

100.0% PROCESSED 592 ITERATIONS 39 ANSWERS
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 10381 TO 13299
PROJECTED ANSWERS: 406 TO 1154

L4 39 SEA SSS SAM L3

=> s sss full l3

FULL SEARCH INITIATED 14:48:03 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 11663 TO ITERATE

100.0% PROCESSED 11663 ITERATIONS 1047 ANSWERS
SEARCH TIME: 00.00.01

L5 1047 SEA SSS FUL L3

=> file caplus medline

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	152.15	152.36

FILE 'CAPLUS' ENTERED AT 14:48:28 ON 13 JUN 2003
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'MEDLINE' ENTERED AT 14:48:28 ON 13 JUN 2003

=> s l5

L6 199 L5

=> s l6 and kappa

L7 113 L6 AND KAPPA

=> s 17 and py<= 1996

L8 58 L7 AND PY<= 1996

=> s 18 and pruritus

L9 0 L8 AND PRURITUS

=> duplicate remove l8

DUPLICATE PREFERENCE IS 'CAPLUS, MEDLINE'

KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n

PROCESSING COMPLETED FOR L8


L10 44 DUPLICATE REMOVE L8 (14 DUPLICATES REMOVED)

L10 ANSWER 2 OF 44 CAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER: 1996:410432 CAPLUS
 DOCUMENT NUMBER: 125:86487
 TITLE: Preparation of N-[2-(1-pyrrolidiny)-1-phenylethyl]acetamide as **kappa**-receptor agonists
 INVENTOR(S): Ito, Fumitaka
 PATENT ASSIGNEE(S): Pfizer Pharmaceuticals Inc., Japan
 SOURCE: PCT Int. Appl., 41 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9606078	A1	19960229	WO 1994-JP1399	19940824 <--
W: JP				
CA 2196885	AA	19960229	CA 1995-2196885	19950518 <--
CA 2196885	C	20010123		
WO 9606077	A1	19960229	WO 1995-IB374	19950518 <--
W: AU, CA, CN, CZ, FI, HU, JP, KR, MX, NO, NZ, PL, RU, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
AU 9523506	A1	19960314	AU 1995-23506	19950518 <--
EP 777649	A1	19970611	EP 1995-917437	19950518
EP 777649	B1	19990714		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
JP 09510731	T2	19971028	JP 1995-529926	19950518
JP 2935899	B2	19990816		
AT 182138	E	19990715	AT 1995-917437	19950518
ES 2133767	T3	19990916	ES 1995-917437	19950518
BR 9503775	A	19960416	BR 1995-3775	19950823 <--
FI 9700746	A	19970221	FI 1997-746	19970221
US 5837720	A	19981117	US 1997-793225	19970417
PRIORITY APPLN. INFO.:			WO 1994-JP1399	A 19940824
			WO 1995-IB374	W 19950518

OTHER SOURCE(S): MARPAT 125:86487
 GI



AB The title compds. [I; R = hydrogen or hydroxy; A = (un)substituted Ph; X = (un)substituted Ph or heterocyclic, mono-, di- or tri-halomethyl, cyano, etc.; X1 = Ph, furyl, thienyl, pyridyl, thiazolyl, benzofuryl, benzothienyl, etc.], which have agonist activity toward opioid **kappa** receptors (no data) and are useful as analgesics (no data), antiinflammatory agents (no data), diuretics (no data), and neuroprotective agents (no data), are prepd. Thus, (2S,3S)-1-[2-N-(benzyloxycarbonyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine was condensed with 3,4-dichlorophenylacetyl chloride and the free base salified with HCl, producing the hydrochloride salt of N-(benzylcarbonyl)methyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.

L10 ANSWER 15 OF 44 CAPLUS COPYRIGHT 2003 ACS DUPLICATE 4
 ACCESSION NUMBER: 1995:217334 CAPLUS
 DOCUMENT NUMBER: 122:9806
 TITLE: **.kappa.** Opioid Receptor Selective Affinity
 Labels: Electrophilic Benzeneacetamides as **.kappa.**-Selective Opioid Antagonists
 AUTHOR(S): Chang, An-Chih; Takemori, Akira E.; Ojala, William H.;
 Gleason, William B.; Portoghese, Philip S.
 CORPORATE SOURCE: College of Pharmacy, University of Minnesota,
 Minneapolis, MN, 55455, USA
 SOURCE: Journal of Medicinal Chemistry (1994),
 37(26), 4490-8
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB Pyrrolidinyloethylacetamides I [R = 3-, 4-NCS, (E)-NHCOCH:CHCO₂Me] were synthesized as **.kappa.**-selective affinity labels and evaluated for opioid activity. In smooth muscle prepns., the non-electrophilic parent compd. (+)-S-I [R = H] and the affinity labels behaved as **.kappa.** agonists in that they were potently antagonized by norbinaltorphimine (norBNI). In addn. to the high binding affinity and selectivity of I [R = 3-NCS] to **.kappa.** opioid receptors, wash studies have suggested that this involves covalent binding. In the mouse tail-flick assay, I [R = NCS] produced long-lasting antagonism of the antinociceptive effect of the **.kappa.** opioid agonist, ((+)-)-trans-2-(3,4-dichlorophenyl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide ((+)-)-U50,488). In contrast, the non-electrophilic parent compd. (+)-S-I [R = H] and I [R = 3-(E)-NHCOCH:CHCO₂Me] were devoid of antagonist activity in the tail-flick assay. At substantially different doses, I [R = 3-, 4-NCS] also produced antinociception in the mouse abdominal stretch assay. In addn., I [R = 3-NCS, 3-(E)-NHCOCH:CHCO₂Me] had improved in vivo **.kappa.** selectivities compared to (+)-S-I [R = H] and I [R = 4-(E)-NHCOCH:CHCO₂Me]. The improved **.kappa.** selectivities of I [R = 3-NCS, 3-(E)-NHCOCH:CHCO₂Me] and the different agonist and antagonist potencies of I [R = 3-, 4-NCS] may be explained resp. by the existence of multiple **.kappa.** agonist binding sites and distinct agonist and antagonist binding sites. In view of the antagonist selectivity and the apparent irreversible binding of I [R = 3-NCS] to **.kappa.** receptors, it may serve as a useful pharmacol. or biochem. tool to investigate **.kappa.** opioid receptors.